

Stable Irregular Dynamics in Complex Neural Networks

Sven Jahnke¹⁻³, Raoul-Martin Memmesheimer¹⁻³ and Marc Timme^{1,2}

¹*Network Dynamics Group, Max-Planck-Institute
for Dynamics & Self-Organization (MPIDS),*

²*Bernstein Center for Computational Neuroscience
(BCCN), 37073 Göttingen, Germany, and*

³*Fakultät für Physik, Georg-August-Universität Göttingen*

(Dated: Fri May 18 13:19:03 CEST 2006)

Abstract

For infinitely large sparse networks of spiking neurons mean field theory shows that a balanced state of highly irregular activity arises under various conditions. Here we analytically investigate the microscopic irregular dynamics in finite networks of arbitrary connectivity, keeping track of all individual spike times. For delayed, purely inhibitory interactions we demonstrate that the irregular dynamics is not chaotic but rather stable and convergent towards periodic orbits. Moreover, every generic periodic orbit of these dynamical systems is stable. These results highlight that chaotic and stable dynamics are equally capable of generating irregular activity.

PACS numbers: 05.45.-a, 89.75.-k, 87.10.+e

Highly irregular dynamics is a prominent feature of multi-dimensional complex systems and often attributed to chaos [1, 2]. Networks of spiking neurons, which interact by sending and receiving electrical pulses (spikes) exhibit very irregular dynamics for a wide range of conditions [3, 4, 5, 6, 7, 8]. For instance, networks of sparse random connectivity may display a balanced state [3] in which excitatory (positive) and inhibitory (negative) inputs to each neuron balance on average and only the fluctuations create spikes at irregular points in time [3, 4]. Their dynamics resemble random (e.g. Poisson) processes with low correlations across different neurons. Mean field theory [4] shows that such irregular balanced activity occurs in networks with excitatory and inhibitory recurrent feedback as well as in networks that receive external excitatory inputs and exhibit recurrent inhibition only [4, 5]. Interestingly, for inhibitory networks recent work [9, 10] suggests that this irregular dynamics might be stable, at least in globally coupled networks and in networks without delay. In particular, Zillmer et al. [10] numerically measured a negative maximal Lyapunov exponent of the irregular, seemingly chaotic spiking dynamics.

In this Letter, we analytically investigate the microscopic dynamics in finite networks of spiking neurons with delayed inhibitory interactions [11]. For networks of arbitrary connectivity we show that almost all spike sequences generated by the network – trajectories of the dynamical system – are stable, i.e. sufficiently small perturbations of such trajectories do not grow with time. In particular, every generic periodic orbit in such inhibitory networks is stable. Moreover, the analysis indicates that all attractors in such networks are stable periodic orbits, which, however, may only be reached after very long and irregular, but also stable transients.

Consider N neurons that interact on a directed network by sending and receiving spikes. The subthreshold dynamics of the neurons' membrane potentials $V_i(t)$ at time t are given by

$$\frac{d}{dt}V_i = f_i(V_i) + \sum_{j=1}^N \sum_{k \in \mathbb{Z}} \varepsilon_{i,j} \delta(t - t_{j,k}^s - \tau_{i,j}) \quad (1)$$

where a smooth function f_i specifies the internal dynamics, $\varepsilon_{i,j} \leq 0$ is the inhibitory coupling strength and $\tau_{i,j} > 0$ the delay time of a synaptic interaction from neuron j to neuron i , and $t_{j,k}^s$ determines the time of the k th spike sent by neuron j . Here $\delta(\cdot)$ is the Dirac delta-function. If a neuron j reaches the threshold potential, $V_j(t^-) = V_{\Theta,j}$, it generates a spike at $t =: t_{j,k}^s$ for some k and is reset, $V_j(t_{j,k}^s) = 0$. We here require that the f_j for all j satisfy

$f_j(V_j) > 0$ and $f_j'(V_j) < 0$ for all $V_j \leq V_{\Theta,j}$ such that in isolation each neuron exhibits oscillatory dynamics.

The neurons are equivalently described [12] by a phase-like variable $\phi_j(t) \in (-\infty, \phi_{\Theta,j}]$ satisfying the linear differential equation

$$d\phi_j/dt = 1 \quad (2)$$

at all non-event times. Upon reaching a phase threshold $\phi_j(t_{j,k}^s) = \phi_{\Theta,j}$, this phase is reset, $\phi_j(t_{j,k}^s) := 0$ and a spike is generated. After a delay time $\tau_{i,j}$ that spike is received by postsynaptic neuron i of neuron j and its phase changes according to

$$\phi_i(t_{j,k}^s + \tau_{i,j}) = H_{\varepsilon_{i,j}}^{(i)} \left(\phi_i \left((t_{j,k}^s + \tau_{i,j})^- \right) \right). \quad (3)$$

This interaction is mediated by the transfer function

$$H_{\varepsilon}^{(i)}(\phi) = U_i^{-1} [U_i(\phi) + \varepsilon] \quad (4)$$

where $U_i(t)$ is the free (all $\varepsilon_{i,j} = 0$) solution of (1) through the initial condition $U_i(0) = 0$, yielding $U_i' > 0$ and $U_i'' < 0$, and $\phi_{\Theta,j} = U_j^{-1}(V_{\Theta,j})$, cf. [13]. For instance, for standard leaky integrate-and-fire neurons, where $f_i(V) = I_i - \gamma_i V$ with time scale $\gamma_i^{-1} \geq 0$ and equilibrium potential $\gamma_i^{-1} I_i > V_{\Theta,i}$, we have $U_i(\phi) = \gamma_i^{-1} I_i (1 - \exp(-\gamma_i \phi))$. Whereas the analysis below is valid for general $U_i(\phi)$, all numerical simulations are presented for integrate-and-fire neurons. In the following we consider arbitrary generic spike sequences in which all neurons are active (i.e. there is a finite $T > 0$, arbitrarily large, such that in every time interval $[t, t + T)$, $t \in \mathbb{R}$, every neuron fires at least once) and no two events occur at the same time. Sending and receiving of spikes are the only nonlinear events occurring in these systems.

Highly irregular spiking sequences (cf. Fig 1a) constitute a typical form of activity in these networks, suggesting that the underlying dynamics may be chaotic. However, as we show below for networks of arbitrary connectivity, this dynamics generically is stable. To show this, we first analytically study the exact microscopic dynamics of an original trajectory, as defined by the (arbitrarily irregular) sequence of events generated by the network, and a slight perturbation to it that keeps the order of events as in the original.

The time of the n th event (sending and receiving) occurring in the entire network is denoted by t_n in the original sequence, and by $\tilde{t}_n$ in the perturbed sequence. Here simultaneous reception of the same spike at different neurons constitute one event. Analogously, at




Figure 1: (color). Stable irregular dynamics in a random network ($N = 400$, $\gamma_i^{-1} I_i \equiv 4.0$, $V_{\Theta,i} \equiv 1$, $\tau_{i,j} \equiv 0.1$, connection probability $p = 0.2$, $\sum_j \varepsilon_{i,j} = -16$). (a) The upper panel displays the spiking times (blue lines) of the first 50 neurons. The lower panel displays the membrane potential trajectory of neuron $i = 1$ (spikes of height $\Delta U = 2$ added at firing times). The inset shows a histogram of the coefficients of variation $CV_i := \sigma_i / \mu_i$; $\mu_i = \langle t_{i,k+1}^s - t_{i,k}^s \rangle$; $\sigma_i^2 := \langle (t_{i,k+1}^s - t_{i,k}^s - \mu_i)^2 \rangle$ averaged over time. (b) Exponential decay of the maximal perturbation $\max_i |\delta_i^{(n)}|$ (blue dots) and the minimal margin $\kappa^{(n)}$ (gray line) for one given microscopic dynamics. (c) Algebraic decay of the average minimal margin, $\overline{\kappa^{(n)}}$ (green dashed line, averaged over 250 random initial conditions) and the analytical prediction (no free fit parameter) of $\overline{\kappa^{(n)}}$ (black solid line). Additionally we show the minimal margin $\kappa^{(n)}$ for three exemplary initial conditions (gray lines), including that of panel b.

a given time t , we denote the phases of neuron i by $\phi_i(t)$ and $\tilde{\phi}_i(t)$, respectively. Let

$$\Delta_i^{(n)} = \left(\phi_i(t_n) - \tilde{\phi}_i(\tilde{t}_n) \right) - (t_n - \tilde{t}_n) := \delta_i^{(n)} - \delta t^{(n)} \quad (5)$$

denote the difference of the phases of neuron i between the two sequences after the n th and before the $(n+1)$ st event, corrected for the time shift $\delta t^{(n)} = t_n - \tilde{t}_n$ between the sequences.

If at the $(n+1)$ st event some neuron j^* sends a spike, the phase shifts

$$\Delta_i^{(n+1)} = \Delta_i^{(n)} \quad (6)$$

of all neurons i stay unchanged. Due to the linear phase dynamics (2) between the spikes, $\Delta_{j^*}^{(n+1)} = \Delta_{j^*}^{(n)} = -\delta t^{(n+1)}$ also specifies the temporal shift of the $(n+1)$ st event. At some $t_l = t_{n+1} + \tau_{i,j^*}$ postsynaptic neuron i receives the spike sent by j^* . The resulting phase shifts are given by

$$\Delta_i^{(l)} = H_{\varepsilon_{i,j^*}}^{(i)}(\phi_i(t_l^-)) - H_{\varepsilon_{i,j^*}}^{(i)}(\tilde{\phi}_i(\tilde{t}_l^-)) - \delta t^{(l)}, \quad (7)$$

where $\phi_i(t_l^-) = \phi_i(t_{l-1}) + t_{n+1} + \tau_{i,j^*} - t_{l-1}$ and $\tilde{\phi}_i(\tilde{t}_l^-) = \tilde{\phi}_i(\tilde{t}_{l-1}) + \tilde{t}_{n+1} + \tau_{i,j^*} - \tilde{t}_{l-1}$ are the phases just before spike reception. Using the identities $\phi_i(t_l^-) = \tilde{\phi}_i(\tilde{t}_l^-) + \Delta_i^{(l-1)} + \delta t^{(n+1)}$ and $\delta t^{(l)} = \delta t^{(n+1)} = -\Delta_{j^*}^{(n)}$, we apply the mean value theorem in eq. (7) and obtain

$$\Delta_i^{(l)} = c_i^{(l)} \cdot \Delta_i^{(l-1)} + (1 - c_i^{(l)}) \cdot \Delta_{j^*}^{(n)} \quad (8)$$

where $c_i^{(l)}$ is given by the derivative $c_i^{(l)} = dH_{\varepsilon_{i,j^*}}^{(i)}(\phi)/d\phi$ for some ϕ between $\phi_i(t_l^-)$ and $\tilde{\phi}_i(\tilde{t}_l^-)$. If neuron j^* is not connected to neuron i , $\varepsilon_{i,j^*} = 0$, the function $H_{\varepsilon_{i,j^*}}^{(i)}(\phi) = H_0^{(i)}(\phi) = \phi$ is the identity map, such that the phase shift stays unchanged, $\Delta_i^{(l)} = \Delta_i^{(l-1)}$; indeed $c_i^{(l)} = dH_0^{(i)}(\phi)/d\phi = 1$ is independent of ϕ . If neuron j^* is connected to i we find $c_i^{(l)}$ bounded by

$$c_{\min} := \min_{\phi,k} \{(H_{\varepsilon_{\min}}^{(k)})'(\phi)\} \leq c_i^{(l)} \leq \max_{\phi,k} \{(H_{\varepsilon_{\max}}^{(k)})'(\phi)\} =: c_{\max} \quad (9)$$

where $\varepsilon_{\max} = \max_{i,j:\varepsilon_{i,j} \neq 0} \{\varepsilon_{i,j}\}$ and $\varepsilon_{\min} = \min_{i,j:\varepsilon_{i,j} \neq 0} \{\varepsilon_{i,j}\}$. The phase is confined to some finite interval $\phi \in [\phi_{\min}, \phi_{\max}]$ which depends on the network parameters. Given that $dH_{\varepsilon}^{(i)}(\phi)/d\phi = U_i'(\phi)/U_i'(U_i^{-1}[U_i(\phi) + \varepsilon])$ and using the monotonicity $U_i' > 0$ and concavity $U_i'' < 0$, we find $c_{\min} > 0$ and $c_{\max} < 1$, independent of the sequence and of the network realization (including its connectivity). Thus the phase shift after receiving a spike is a weighted average of earlier shifts.

Consider that the perturbed sequence is created from the original one by perturbing the phases of all neurons and the sending times of all spikes sent but not received at time $t = t_0$. (We denote the maximum of these perturbations by $\Delta_{\max}^{(0)}$ and the minimum by $\Delta_{\min}^{(0)}$). After some finite time all spikes perturbed initially arrived and all perturbations to newly generated spike sending times stem from previous perturbations of some neurons' phases

[14]. Given that the maximum phase shift cannot increase and the minimum not decrease according to (8) and (9), this implies Lyapunov stability of the dynamical trajectories.

For strongly connected networks [15], more involved graph theoretical arguments show that the initial perturbations actually decay exponentially such that the considered trajectories are asymptotically stable (cf. [5, 16]). Briefly, following the propagation of a phase perturbation of one specific neuron l_0 across the entire network, shows that after a finite number of $K := 2NM$ events all perturbations (of the phases of all neurons and of the sending times of spikes sent but not yet received) are bounded from above by

$$\Delta_{\text{mx}}^{(K)} \leq c^* \Delta_{l_0}^{(0)} + (1 - c^*) \Delta_{\text{mx}}^{(0)}. \quad (10)$$

Here $c^* := (1 - c_{\text{mx}})^N (c_{\text{mn}})^{2NM} \leq (1 - c_{\text{mx}})(c_{\text{mx}})$ such that $3/4 \leq (1 - c^*) < 1$ and M depends on the spike sequence but is finite. Similarly we find a lower bound given by $\Delta_{\text{mn}}^{(K)} \geq c^* \Delta_{l_0}^{(0)} + (1 - c^*) \Delta_{\text{mn}}^{(0)}$. The difference of the maximal and the minimal perturbation after K events is therefore given by

$$\delta_i^{(K)} \leq \Delta_{\text{mx}}^{(K)} - \Delta_{\text{mn}}^{(K)} \leq (1 - c^*)(\Delta_{\text{mx}}^{(0)} - \Delta_{\text{mn}}^{(0)}). \quad (11)$$

This inequality (11) implies that both sequences converge exponentially fast against each other. Thus all sequences considered are asymptotically stable for all strongly connected networks.

A main condition for stability of trajectories was that the order of events stays the same in the perturbed and original trajectories. For arbitrary non-degenerate spike sequences, there is a non-zero perturbation size keeping the order unchanged in any *finite* time interval. However, the requirement of an unchanged event order yields more and more conditions over time such that the allowed size of a perturbation could decay more quickly with time than the actual perturbation. This will be excluded if the temporal *margin* $\mu^{(n)}$ between two subsequent events stays larger than the dynamical perturbation for *infinite* time. Formally, after time t_n denote the k th potential future event time (of the original trajectory) that would arise if there were no future interactions by $\theta_{n,k}$, $k \in \mathbb{N}$, and the temporal *margin* by $\mu^{(n)} := \theta_{n,2} - \theta_{n,1}$. A sufficiently small perturbation, satisfying $\Delta_{\text{mx}}^{(n)} - \Delta_{\text{mn}}^{(n)} \leq \mu^{(n)}$, cannot change the order of the $(n + 1)$ st event.

This directly implies that almost all *periodic orbits* (all those with non-degenerate event times t_n) consisting of a finite number of P events are stable because there is a minimal

margin

$$\kappa^{(P)} := \min_{n \in \{1, \dots, P\}} \mu^{(n)} \quad (12)$$

for every non-degenerate periodic pattern.

To further analyze stability properties of *irregular non-periodic spike sequences*, we consider the minimal margin $\kappa^{(n)}$ over the first n events. Assuming that, along with the irregular dynamics, the temporal margins are generated by a Poisson point process with rate ν , the distribution function of margins is given by $P(\mu^{(n)} \leq \mu) = 1 - e^{-\nu\mu}$. The probability that the minimal margin $\kappa^{(n)}$ after n events is smaller or equal to μ is determined by the probabilities that not all individual margins $\mu^{(n)}$ are larger than μ such that

$$P(\kappa^{(n)} \leq \mu) = 1 - \prod_{m=1}^n P(\mu^{(m)} > \mu) = 1 - e^{-n\nu\mu} \quad (13)$$

with density $\rho_n(\mu) := dP(\kappa^{(n)} \leq \mu)/d\mu = n\nu \exp(-n\nu\mu)$. This implies an algebraic decay with the number n of events for the expected minimal margin

$$\overline{\kappa^{(n)}} = \int_0^\infty \mu \rho_n(\mu) d\mu = (\nu \cdot n)^{-1} \quad (14)$$

that depends only on the event rate and is independent of the specific network parameters. Numerical simulations confirm this *algebraic* decay (14) of the expected minimal margin with the number of network events n ; an example is shown in Fig. 1c. Together with the *exponential* decay of dynamical perturbations (11) this indicates that for a sufficiently small perturbation the order of events stays unchanged for *infinite* time.

Interestingly, arbitrary irregular spike sequences necessarily converge to a periodic orbit after finite time (cf. Fig. 2) because there is some finite number E such that (i) two sequences that share the same order of E events are equally ordered for all future events because any initial difference decays exponentially (11) and (ii) there is only a finite number of orderings of events in a finite network such that a given sequence of length E repeats after finite time. Nevertheless, we find that the transient time until a periodic orbit is reached, rapidly increases with network size N and with the interaction strengths, in agreement with observations in ref. [10]; cf. also [16].

In summary, we analytically accessed the microscopic dynamics of inhibitory networks of spiking neurons with arbitrarily complicated connectivity and delayed interactions. We showed that all generic trajectories are stable, even if they generate highly irregular dynamics. In particular, the results analytically confirm recent numerical findings [10] that irregular




Figure 2: (color) Convergence towards a periodic orbit in a random network ($N = 40$, $\gamma_i^{-1} I_i \equiv 3.0$, $V_{\Theta,i} \equiv 1.0$, $\tau_{i,j} \equiv 0.1$, $p = 0.2$, $\sum_j \epsilon_{i,j} = -3.3$). (a) The average minimal margin $\overline{\kappa}^{(n)}$ (as in Fig. 1c) decays as a power-law (region A) and saturates after about 10^7 events (region B) when the periodic orbit is reached. Inset: Margin $\mu^{(n)}$ (black) and minimal margin $\kappa^{(n)}$ (gray) for a trajectory started from one specific initial condition. The margin $\mu^{(n)}$ fluctuates strongly on the transient but is comparatively large and bounded after the sequence becomes periodic; thus the minimal margin $\kappa^{(n)}$ does not decrease further for future events n . (b), (c): Snapshots of irregular spike sequences (b) after $n = 20.000$ events on the transient and (c) after $n = 10^8$ events on the periodic orbit.

dynamics in spiking neural networks may exhibit stable behavior. Moreover, as for globally coupled systems without delay [9] our results show that also networks with more complex structure and delayed interactions exhibit stable periodic orbits. However, highly irregular yet stable transient trajectories dominate the dynamics in all large and non-globally coupled networks, in stark contrast to the fast convergence to attractor dynamics found recently in globally coupled networks [9] and also opposite to the long chaotic (and thus unstable) transients in randomly diluted networks of excitatory neurons [6]. In particular, in the class of systems presented here, almost every trajectory is stable, not only if it generates regular [5] but even if it generates very irregular dynamics. Curiously, the assumption of events generated by a maximally irregular (Poisson) random process led us to show stability of the deterministic trajectories.

More generally, our results underline that multi-dimensional deterministic dynamics with statistical properties close to that of a random system need not be generated by deterministic

chaos. As we have shown analytically for spiking neural networks with delayed interactions, dynamical irregularity may well be generated by stable trajectories, dominantly on a transient not on an attractor. Future work needs to investigate closer the key consequences for systems in which irregular dynamics is stable. For instance stable irregular dynamics lifts the important practical constraint of long-term unpredictability that irregular dynamics bears if it is generated by chaos. Stable irregular dynamics, even in multi-dimensional systems, may well be predictable in practice. For the networks of spiking neurons studied above, this has the astounding consequence that the dynamics in only a small time window, even in the presence of some errors, defines the entire future of the highly irregular spiking dynamics.

We thank Fred Wolf for helpful comments and the Federal Ministry of Education and Research (BMBF) Germany for partial support under Grant No. 01GQ0430.

-
- [1] R. Badii and A. Politi, *Complexity* (Cambridge University Press, 1999).
 - [2] P. Cvitanović, R. Artuso, R. Mainieri, G. Tanner, and G. Vattay, *Chaos: Classical and Quantum* (ChaosBook.org, Niels Bohr Institute, Copenhagen, 2005).
 - [3] C. v. Vreeswijk and H. Sompolinsky, *Science* **274**, 1724 (1996); *Neural Comput.* **10**, 1321 (1998).
 - [4] N. Brunel and V. Hakim, *Neural Comput.* **11**, 1621 (1999); N. Brunel, *J. Comput. Neurosci.* **8**, 183 (2000).
 - [5] M. Timme, F. Wolf, and T. Geisel, *Phys. Rev. Lett.* **89**, 154105 (2002); M. Timme and F. Wolf, in prep.
 - [6] A. Zumdick, M. Timme, T. Geisel, and F. Wolf, *Phys. Rev. Lett.* **93**, 244103 (2004).
 - [7] D. Hansel and G. Mato, *Phys. Rev. Lett.* **86**, 4175 (2001).
 - [8] C. v. Vreeswijk, *Phys. Rev. Lett.* **84**, 5110 (2000).
 - [9] D. Jin, *Phys. Rev. Lett.* **89**, 208102 (2002).
 - [10] R. Zillmer, R. Livi, A. Politi, and A. Torcini, *Phys. Rev. E* **74**, 036203 (2006).
 - [11] Delays make the system formally infinite dimensional; for the systems considered here, however, the dimension is generally high but finite [14].
 - [12] R. Mirollo and S. Strogatz, *SIAM J. Appl. Math.* **50**, 1645 (1990); U. Ernst, K. Pawelzik, and T. Geisel, *Phys. Rev. Lett.* **74**, 1570 (1995).

- [13] R. Memmesheimer and M. Timme, *Physica D* **224**, 182 (2006).
- [14] P. Ashwin and M. Timme, *Nonlinearity* **18**, 2035 (2005).
- [15] A network is *strongly connected* if there is a directed path of connections between any ordered pair of neurons.
- [16] S. Jahnke, R. Memmesheimer, and M. Timme, in prep.